

THE DANGER OF INDUSTRIALIZATION

AIR POLLUTION IN ALASKA'S NORTH SLOPE

And its implications for the community of Nuiqsut

By Vetta Stepanyan | February 2019

This project was completed as part of an internship with Alaska Community Action on Toxics for graduate credit through the National University of Natural Medicine.



1225 E. International Airport Road; Suite 220
Anchorage, Alaska 99518-1410

www.akaction.org

Telephone: (907) 222-7714 | Fax: (907) 222-7715

Introduction

Alaska's North Slope is the region north of the crest of the Brooks Range and is one of the most extreme environments in which humans live. Summer temperatures range between 40-60°F, while winter temperatures sometimes fall below -40°F.²⁸

There are eight communities in Alaska's North Slope: Anaktuvuk Pass, Atkasuk, Barrow, Kaktovik, Nuiqsut, Point Hope, Point Lay, and Wainwright. The community of Nuiqsut is located 136 miles southeast of Barrow, the northernmost city in the United States of America.

According to the U.S. Census, there were 402 residents living in the community of Nuiqsut in 2010. Nuiqsut's economy is based on subsistence hunting, fishing, and whaling and local knowledge encompasses comprehensive information about various aspects of the environment.

Massive oil reserves were discovered in Alaska's North Slope in 1968.²⁸ Following that initial discovery, oil and gas facilities have continued their expansion throughout this region. As a result of this continued expansion, air pollution is of increasing concern to residents of Nuiqsut which was first surrounded by an oil drilling unit in 2000 with a subsequent cluster of childhood leukemia and persistent asthma in the children which were reported by Nuiqsut leaders to ACAT staff in 2012. By 2018 the oil rigs had surrounded the village. The burden of the socioeconomic, environmental, and health costs of oil exploration, development, and production has fallen upon them.

Considerable research has been done on the effects of oil and gas activities, but information on the cumulative effects on human health is lacking, especially in the Arctic region. Furthermore there needs to be a focus on the impacts of oil and gas activities on the health and wellbeing of individuals and their communities in the North Slope as it relates to their traditional ways of life.

In this report we will present the inherent risks that oil and gas activities have on environmental health by calling attention to the hazardous air pollutants that are released into the ecosystem on Alaska's North Slope.

Methods

For this report, ACAT compiled all emissions data using the 'facility mapping' tool found on The National Emissions Inventory (NEI) webpage, which is published by the US Environmental Protection Agency (EPA). The inventory is released every three years, and we used data that were reported in 2008, 2011, and 2014 that represent emissions released over the previous three-years, respectively.

The inventory provides a detailed and comprehensive estimate of the criteria air pollutants and hazardous air pollutants that have been released from various industrial sources throughout Alaska. We narrowed our research to include only those facilities that were constituted under the North American Industry Classification System (NAICS) as: Support Activities for Oil and Gas Operations, Pipeline Transportation of Crude Oil, Crude Petroleum and Natural Gas Extraction, or Natural Gas Distribution. Furthermore, we focused only on those facilities that are in Alaska's North Slope Borough. ACAT compiled these data and highlighted the top pollutants from each inventory year. The NEI webpage offered some emissions quantities in tons and others in pounds. To make the data homogenous, we converted tons into pounds for all relevant pollutants.


ACAT used PubMed to research the harmful health impacts of the most frequent pollutants reported in the NEI. With this information, we created a health impact table to help guide the evaluation of the health hazards posed by these air pollutants that were released of the nine years covered by the NEI and continue to be released into the air on Alaska's North Slope.

To learn more about the National Emissions Inventory, please visit:

<https://www.epa.gov/air-emissions-inventories>

For all emissions inventories (years 2008, 2011, and 2014), Nitrogen Oxides comprised the largest emissions.


ACAT Graph 1. Nitrogen Oxides Emissions Over Time



The graph below highlights the magnitude of change in the amount of Nitrogen Oxides emissions over the three inventory years.

Although there is an observed decrease in the amount of Nitrogen Oxides emissions over time, please draw your attention to the total value of emissions across the trend, from 2008 to 2014. Over 200,000,000 pounds of Nitrogen Oxides have been emitted over a 9-year period (from 2006 through 2014).

ACAT Graph 2. Emissions Trends for Nitrogen Oxides




Emissions Inventory Year: 2008

Top pollutants in pounds: Sulfur Dioxides, PM10, PM2.5, VOCs, Acetaldehyde, Benzene, Ethyl Benzene, Formaldehyde, Toluene, and Xylenes.

ACAT Table 1: Top 11 Pollutants (in Pounds) for 2008

Air Pollutant	Emissions (in pounds)
Nitrogen Oxides	72,523,580
Sulfur Dioxides	1,859,760
PM10	1,809,440
PM2.5	1,785,497.08
VOC	1,478,720.00
Acetaldehyde	10,711.59
Benzene	3,115.19
Ethyl Benzene	8,045.18
Formaldehyde	183,933.31
Toluene	29,584.32
Xylenes	14,561.69

ACAT Graph 3. Top Hazardous Air Emission (2008)




Emissions Inventory Year: 2011

Top pollutants in pounds: Sulfur Dioxides, PM10, PM2.5, VOCs, Acetaldehyde, Benzene, Ethyl Benzene, Formaldehyde, Toluene, and Xylenes.

ACAT Table 2: Top Pollutants (in Pounds) For 2011

Air Pollutant	Emissions (in pounds)
Nitrogen Oxides	72,382,940
Sulfur Dioxide	1,878,680
PM10	1,847,020
PM2.5	1,810,839.48
VOC	978,560.00
Acetaldehyde	8,794.90
Benzene	2,478.82
Ethyl Benzene	6,603.14
Formaldehyde	168,848.18
Toluene	28,813.54
Xylenes	13,769.36

ACAT Graph 4: Top Hazardous Air Emissions (2011)




Emissions Inventory Year: 2014

Top pollutants in pounds: Sulfur Dioxides, PM10, PM2.5, VOCs, Acetaldehyde, Hexane, Phenol, Formaldehyde, Toluene, and Xylenes.

ACAT Table 3: Top Pollutants (in Pounds) for 2014


Air Pollutant	Emissions (in pounds)
Nitrogen Oxides	68,695,420
Sulfur Dioxides	2,184,980
PM10	1,776,668
PM2.5	1,733,795.19
VOC	1,343,440.00
Acetaldehyde	1,790.26
Hexane	4,402.28
Phenol	2,194.84
Formaldehyde	35,376.25
Toluene	5,105.55
Xylenes	2,509.02

ACAT Graph 5. Top Hazardous Air Emissions (2014)




Sulfur Dioxide comprised the second largest emissions for all three inventory years (2008, 2011, and 2014).

ACAT Graph 6. Emissions Trends for Sulfur Dioxide




The seven graphs below represent emissions trends for the other major pollutants.


ACAT Graph 7. Emissions Trends for PM10




ACAT Graph 8. Emissions Trends for PM2.5




ACAT Graph 9. Emissions Trends for VOCs




ACAT Graph 10. Emissions Trends for Acetaldehyde




ACAT Graph 11. Emissions Trends for Formaldehyde



ACAT Graph 12. Emissions Trends for Toluene



ACAT Graph 13. Emissions Trends for Xylenes



ACAT Table 4. Summary of Adverse Health Impacts for Major Harmful Air Pollutants

Pollutant	Health Impacts
Nitrogen Oxides	NO ₂ specifically: lower logical memory ¹ , more severe allergic responses, reduced pulmonary function, asthma, lower birth weight, and increased risk of preterm birth ² , increased risk of stroke ⁴ , deteriorates spatial learning and potentiates amyloid production ⁷ , contributes to an increased incidence of chronic cough ²⁰
Sulfur Dioxides	Respiratory irritant ²⁰ , increased risk of stroke ⁴ , induces inflammation of membranes, causes bronchial narrowing, and slows mucus flow ²³
PM10	Contributes to more severe allergic responses, increased risk of preterm birth, increased risk of pneumonia, and reduced lung function ²
PM2.5	Penetrates deep in to the respiratory tract wherein it can be absorbed in to the blood stream ²¹ , can be translocated to organ tissue through blood circulation, contributes to more severe allergic responses, decreased birth weight, and asthma ² , lower verbal learning performance ¹ , increased cardiovascular mortality ^{4,5} , reduced cardiovascular function ^{5,22} , has the ability to enter the olfactory epithelium and can be transported to the olfactory bulb causing olfactory dysfunction ¹⁹ , induces inflammatory reactions across organ systems ²²
Formaldehyde	Eye, nose, and throat irritant resulting in cough, wheezing, chest pains, and bronchitis; a carcinogen resulting in increased incidence of lung and nasopharyngeal cancer ^{5, 26}
Toluene	Physiological depression of the central nervous system ^{16,17} , cardiotoxic ¹⁷ , causes renal tubular acidosis and can cause headache, dizziness, confusion, muscle weakness, and even muscle paralysis ¹⁷
Xylenes	Nose and throat irritation, severe lung congestion, pulmonary hemorrhages, edema, impaired short-term memory, as well as alteration in equilibrium or body balance ⁸ , reduced muscle power, depression of the central nervous system inducing symptoms such as headache, dizziness, and vomiting ⁹ , pathological changes in ovarian tissue, ovary atrophy ¹⁰
Acetaldehyde	Carcinogenic and genotoxic ^{11, 27} , can cause mild respiratory irritation ²⁷
Ethyl Benzene	Ototoxic (having a toxic effect on the ear or its nerve supply) ^{24, 25}
Benzene	Reduced pulmonary function ² , decreases the number of cells in bone marrow causing blood disorders ^{2,5} , genotoxic causing genetic damage including DNA cross linking and sister chromatid exchanges ^{3,18} , increases cardiovascular risk and injury ⁶ , shortness of breath and lethargy ¹⁷ , carcinogenic ¹⁸
Phenol	Accelerates pubertal development and disrupts estrogenic activity ^{12, 13}
Hexane	Inhibits follicular development, damages ovarian cell ultrastructure, and can cause menstrual abnormalities ¹⁴ , gestational inhalation can alter the reproductive cycle of female offspring ¹⁵

References Cited for this Report

1. Gatto N, Henderson VW, Hodis HN, et al. Components of air pollution and cognitive function in middle-aged and older adults in Los Angeles. *NeuroToxicology*. 2014; 40: 1-7.
2. Kim D, Chen Z, Zhou L, Huang S. Air pollutants and early origins of respiratory diseases. *Chronic Dis Transl Med*. 2018; 4(2): 75-94.
3. Duarte-Davidson R, Courage C, Rushton L, Levy L. Benzene in the environment: an assessment of the potential risks to the health of the population. *Occup Environ Med*. 2001; 58(1): 2-13.
4. Bourdrel T, Bind MA, Bejot Y, Morel O, Argacha JF. Cardiovascular effects of air pollution. *Arch Cardiovasc Dis*. 2017; 110(11): 634-642.
5. Suh HH, Bahadori T, Vallarino J, Spengler JD. Criteria air pollutants and toxic air pollutants. *Environ Health Perspect*. 2000; 108(4): 625-633.
6. Abplanalp W, DeJarnett N, Riggs DW, et al. Benzene exposure is associated with cardiovascular risk. *PLoS One*. 2017; 12(9).
7. Yan W, Yun Y, Ku T, Li G, Sang N. NO₂ inhalation promotes Alzheimer's disease-like progression: cyclooxygenase-2-derived prostaglandin E₂ modulation and monoacylglycerol lipase inhibition-targeted medication. *Sci Rep*. 2016; 6: 22429.
8. Reese E, Kimbrough R. Acute toxicity of gasoline and some additives. *Environ Health Perspect*. 1993; 101(6): 115-131.
9. Kandyala R, Raghavendra SPC, Rajasekharan ST. Xylene: an overview of its health hazards and preventive measures. *J Oral Maxillofac Pathol*. 2010; 14(1): 1-5.
10. Niaz K, Bahadar H, Maqbool F, Abdollahi M. A review of environmental and occupational exposure to xylene and its health concerns. *EXCLI J*. 2015; 14:1167-1186.
11. Moeller BC, Recio L, Green A, et al. Biomarkers of exposure and effect in human lymphoblastoid TK6 cells following [¹³C₂]- acetaldehyde exposure. *Toxicol Sci*. 2013; 133(1): 1-12.
12. Wolff MS, Teitelbaum SL, Pinney SM, et al. Investigation of relationships between urinary biomarkers of phytoestrogens, phthalates, and phenols and pubertal stages in girls. *Environ Health Perspect*. 2010; 118(7): 1039-1046.

13. Rasier G, Toppari J, Parent AS, Bourguignon JP. Female sexual maturation and reproduction after prepubertal exposure to estrogens and endocrine disrupting chemicals: a review of rodent and human data. *Mol Cell Endocrinol*. 2006; 254-255: 187-201.
14. Jin L, Ling HH, Fen P, Chang ZW. The effect of n-hexane on the gonad toxicity of female mice. *Biomed Environ Sci*. 2012; 25(2):189-196.
15. Li H, Zhang C, Ni F, et al. Gestational n-hexane alters the expression of genes related to ovarian hormone production and DNA methylation states in adult female F1 rat offspring. *Toxicol Lett*. (239)3: 141-151.
16. Cohr KH. Toluene. A toxicological review. *Scand J Work Environ Health*. 1979; 5(2): 71-90.
17. Vitale CM, Gutovitz S. Aromatic (Benzene, Toluene) Toxicity. Treasure Island, FL: StatPearls Publishing. [https://www-ncbi-nlm-nih.gov/nunm.idm.oclc.org/books/NBK532257/](https://www.ncbi-nlm-nih.gov/nunm.idm.oclc.org/books/NBK532257/). Published October 27, 2018. Accessed January 23, 2019.
18. Arnold SM, Angerer J, Boogaard PJ, et al. The use of biomonitoring data in exposure and human health risk assessment: benzene case study. *Crit Rev Toxicol*. 2013; 43(2):119-153.
19. Ajmani GS, Suh HH, Pinto JM. Effects of ambient air pollution on olfaction: a review. *Environ Health Perspect*. 2016; 124(11): 1683-1693.
20. Groneberg-Kloft B, Kraus T, Mark A, Wagner U, Fischer A. Analysing the cause of chronic cough: relation to diesel exhaust, ozone, nitrogen dioxides, sulphur oxides, and other environmental factors. *J Occup Med Toxicol*. 2006; 1:6.
21. Traboulsi H, Guerrina N, Iu M, Maysinger D, Ariya P, Baglolle C. Inhaled pollutants: the molecular scene behind respiratory and systemic diseases associated with ultrafine particulate matter. *Int J Mol Sci*. 2017; 18(2): 243.
22. Nemmar A, Holme JA, Rosas I, Schwarze PE, Alfaro-Moreno E. Recent advances in particulate matter and nanoparticle technology: a review of the in vivo and in vitro studies. *Biomed Res Int*. 2013; 2013: 279371.
23. Rall DP. Review of the health effects of sulfur oxides. *Environ Health Perspect*. 1974; 8: 97-121.

- 24 Cappaert NLM, Klis SFL, Muijser H, et al. Ototoxic effects of ethyl benzene in rats. *Hear Res.* 1999; 137: 91-102.
25. Vyskocil A, Leroux T, Truchon G, et al. Ethyl benzene should be considered ototoxic at occupationally relevant exposure concentrations. *Toxicol Ind Health.* 2008; 24(4): 241-246.
26. Conolly RB, Kimbell JS, Janszen D, et al. Human respiratory tract cancer risks of inhaled formaldehyde: dose-response predictions derived from biologically-motivated computational modeling of a combined rodent and human dataset. *Toxicol Sci.* 2004; 82(1): 279-296.
27. Sapkota M, Wyatt TA. Alcohol, aldehydes, adducts, and airways. *Biomolecules.* 2015; 5: 2987-3008.
28. National Research Council. *Cumulative Environmental Effects of Oil and Gas Activities on Alaska's North Slope.* Washington, DC: The National Academies Press; 2003. <https://doi.org/10.17226/10639>.